AWARD NUMBER: W81XWH-15-1-0005

TITLE: Demonstrating the Efficacy of Group Prolonged Exposure Treatment of PTSD in OEF/OIF/OND Male Veterans

PRINCIPAL INVESTIGATOR: K. Janet C'de Baca

CONTRACTING ORGANIZATION: Biomedical Research Institute of New Mexico Albuquerque, NM 87108

REPORT DATE: December 2015

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

# REPORT DOCUMENTATION PAGE

c. THIS PAGE

Unclassified

b. ABSTRACT

Unclassified

a. REPORT

Unclassified

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

A TITLE AND SUBTILEE  Demonstrating the Efficacy of Group Prolonged Exposure Treatment of PTSD in OEF/OIF/OND Male Veterans  5b. GRANT NUMBER MS1XWII-15-1-0005  5c. PROGRAM ELEMENT NUMBER  6. AUTHOR(S) Janet Kathreen C'de Baca  5d. PROJECT NUMBER  5d. WORK UNIT NUMBER  5d. TASK NUMBER  5d. WORK UNIT NUMBER  5d. WORK UNIT NUMBER  15d. San Pedro Drive SE; Bldg. 14 (151) Abouquerque, NM 87108  5d. PROSPANIES OF SE; Bldg. 14 (151) Abouquerque, NM 87108  5d. PROSPANIES OF SE; Bldg. 14 (151) Abouquerque, NM 87108  5d. PROSPANIES OF SE; Bldg. 14 (151) Abouquerque, NM 87108  5d. PROSPANIES OF SE; Bldg. 14 (151) Abouquerque, NM 87108  5d. PROSPANIES OF SE; Bldg. 14 (151) Abouquerque, NM 87108  5d. PROSPANIES OF SE; Bldg. 14 (151) Abouquerque, NM 87108  5d. PROSPANIES OF SE; Bldg. 14 (151) Abouquerque, NM 87108  5d. WORK UNIT NUMBER  5d. LEMENT NUMBER  5d. WORK UNIT NUMBER  5d. LEMENT NUMBER  5d. LEMENT NUMBER  5d. WORK UNIT NUMBER  5d. LEMENT NUMBER 5d. LEMENT NUMBER 5d. LEMENT NUMBER 5d. LEMENT NUMBER 5d. L	1. REPORT DATE	2. REPORT TYPE		3. E	ATES COVERED
Demonstrating the Efficacy of Group Prolonged Exposure Treatment of PTSD in OEF/OIF/OND Male Veterans  5b. GRANT NUMBER		Annual		1	0 Nov 2014 - 9 Nov 2015
OEF/OIF/OND Male Veterans    M811XM1-15-1-0005	4. TITLE AND SUBTITLE			5a.	CONTRACT NUMBER
OEF/OIF/OND Male Veterans  6. AUTHOR(S) Janet Kathreen C'de Baca  5d. PROJECT NUMBER  5d. PROJECT NUMBER  5c. TASK NUMBER  5c					
OEF/OIF/OND Male Veterans  6. AUTHOR(S) Janet Kathreen C'de Baca  5d. PROJECT NUMBER  5d. PROJECT NUMBER  5c. TASK NUMBER  5c	Demonstrating the Efficacy of Group Prolonged Exposure		e Treatment of PTSI	) in 5b.	GRANT NUMBER
56. AUTHOR(S) Janet Kathreen C'de Baca  56. PROJECT NUMBER  56. TASK NUMBER  57. WORK UNIT NUMBER  57. WORK UNIT NUMBER  58. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  88. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  88. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  88. PERFORMING ORGANIZATION REPORT NUMBER  100. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  100. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  101. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  102. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  113. SUPPLEMENTARY NOTES  14. ABSTRACT  This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memory is or stimulations no longer result in anxieous arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment of PTSD, where the mechanisms of change include altering current maladaptive relational patterns of behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS  16. AUTHOR STAN AND ADDRESS(ES)  17. LIMITATION			, modument of the	W8	1XWH-15-1-0005
Sd. PROJECT NUMBER  5d. PROJECT NUMBER  5e. TASK NUMBER  5f. WORK UNIT NUMBER  8f. PEFFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Biomedical Research Institute of New Mexico  1501 San Pedro Drive SE; Bldg. 14  (151) Albuquerque, NM 87108  9f. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSOR/MONITOR'S REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT  This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1, The Prolonged Exposure Group treatment will significantly ower PTSD symptoms compared to the Present-Centered Therapy control rogroup. Hypothesis 2, While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT				50	PROGRAM FLEMENT NUMBER
Janet Kathreen Cde Baca  E-Mail: Kathreen CdeBaca@va.gov 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Biomedical Research Institute of New Mexico 1501 San Pedro Drive SE; Bldg. 14 (151) Albuquerque, NM 87108 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged reposure Group resument will significantly lower PTSD symptoms compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBLECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND				30.	TROOKAM ELEMENT NOMBER
Janet Kathreen Cde Baca  E-Mail: Kathreen CdeBaca@va.gov 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Biomedical Research Institute of New Mexico 1501 San Pedro Drive SE; Bldg. 14 (151) Albuquerque, NM 87108 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged reposure Group resument will significantly lower PTSD symptoms compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBLECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND	C AUTHOR(C)			F-1	DDO IECT NUMBER
E-Mail: Kathreen.CdeBaca@va.gov 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Biomedical Research Institute of New Mexico 1501 San Pedro Drive SE; Bldg. 14 (151) Albuquerque, NM 87108 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012 11. SPONSOR/MONITOR'S ACRONYM(S) 12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited 13. SUPPLEMENTARY NOTES 14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly decrease for subjects in the PEG compared to the Present-Centered Therapy control group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TEMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND				50.	PROJECT NUMBER
E-Mail: Kathreen.CdeBaca@va.gov 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION REPORT NUMBER 8. PERFORMING ORGANIZATION REPORT NUMBER 8. PERFORMING ORGANIZATION REPORT NUMBER 9. SPONSORING 7 MONITORING AGENCY NAME(S) AND ADDRESS(ES) 9. SPONSORING 7 MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORIMO 7 MONITORING AGENCY NAME(S) AND ADDRESS(ES) 11. SPONSOR/MONITOR'S ACRONYM(S) 12. DISTRIBUTION / AVAILABILITY STATEMENT 13. SUPPLEMENTARY NOTES 14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so tituations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship	Janet Kathreen C de Baca				
E-Mail: Kathreen.CdeBaca@va.gov 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION REPORT NUMBER 8. PERFORMING ORGANIZATION REPORT NUMBER 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 11. SPONSOR/MONITOR'S ACRONYM(S) 12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited 13. SUPPLEMENTARY NOTES 14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptomatology, the PEG will show greater withing compared to the PCT group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater withing in provement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND				5e.	TASK NUMBER
E-Mail: Kathreen.CdeBaca@va.gov 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION REPORT NUMBER 8. PERFORMING ORGANIZATION REPORT NUMBER 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 11. SPONSOR/MONITOR'S ACRONYM(S) 12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited 13. SUPPLEMENTARY NOTES 14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptomatology, the PEG will show greater withing compared to the PCT group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater withing in provement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Biomedical Research Institute of New Mexico 1501 San Pedro Drive SE; Bldg. 14 (151) Albuquerque, NM 87108  9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSORIMO/ MONITORIS REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT  This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors. Protologed Exposure Group treatment will significantly lower PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects i				5f. \	WORK UNIT NUMBER
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Biomedical Research Institute of New Mexico 1501 San Pedro Drive SE; Bldg. 14 (151) Albuquerque, NM 87108  9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSORIMO/ MONITORIS REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT  This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors. Protologed Exposure Group treatment will significantly lower PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects i	E-Mail: Kathreen.CdeBaca@va.gov	<b>/</b>			
Biomedical Research Institute of New Mexico 1501 San Pedro Drive SE; Bldg. 14 (151) Albuquerque, NM 87108  9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSORIMONITOR'S REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptomos compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group, Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  16. SECURITY CLASSIFICATION OF: 17. LIMITATION 18. NUMBER 19a. NAME OF RESPONSIBLE PERSON	7. PERFORMING ORGANIZATION NAME(	S) AND ADDRESS(ES)		8. F	ERFORMING ORGANIZATION REPORT
Mexico 1501 San Pedro Drive SE; Bldg. 14 (151) Albuquerque, NM 87108  9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSOR/MONITOR'S ACRONYM(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptomes compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater with		-,			
Mexico 1501 San Pedro Drive SE; Bldg. 14 (151) Albuquerque, NM 87108  9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSOR/MONITOR'S ACRONYM(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptomes compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater with	Biomedical Research Institute of Ne	ew			
1501 San Pedro Drive SE; Bidg. 14 (151)  9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSOR/MONITOR'S REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing syscho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND					
(151) Albuquerque, NM 87108  9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSOR/MONITOR'S REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND		I			
Albuquerque, NM 87108  9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSOR/MONITOR'S REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS  PTSD, Group Treatment, Male Veterans, OEF/OIF/OND		•			
U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSOR/MONITOR'S REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON					
U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012  11. SPONSOR/MONITOR'S REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON					
Tot Detrick, Maryland 21702-5012  11. SPONSOR/MONITOR'S REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT  This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS  PTSD, Group Treatment, Male Veterans, OEF/OIF/OND	9. SPONSORING / MONITORING AGENCY	' NAME(S) AND ADDRES	S(ES)	10.	SPONSOR/MONITOR'S ACRONYM(S)
Tot Detrick, Maryland 21702-5012  11. SPONSOR/MONITOR'S REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT  This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS  PTSD, Group Treatment, Male Veterans, OEF/OIF/OND					
Tot Detrick, Maryland 21702-5012  11. SPONSOR/MONITOR'S REPORT NUMBER(S)  12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT  This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS  PTSD, Group Treatment, Male Veterans, OEF/OIF/OND	U.S. Army Medical Research and N	Materiel Command			
12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT  This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS  PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON				11.	SPONSOR/MONITOR'S REPORT
12. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON	Torr Betriok, Maryland 21702 0012	-			NUMBER(S)
Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND					
Approved for Public Release; Distribution Unlimited  13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND	40 DICTRIBUTION / AVAIL ADJUTY CTAT	FRACAIT			
13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problems solving strategies to address present life problems. <a href="https://hypothesis.1">hypothesis.1</a> . The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. <a href="https://hypothesis.2">hypothesis.2</a> . While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. <a href="https://hypothesis.3">hypothesis.3</a> . Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON	12. DISTRIBUTION / AVAILABILITY STAT	EMENI			
13. SUPPLEMENTARY NOTES  14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problems solving strategies to address present life problems. <a href="https://hypothesis.1">hypothesis.1</a> . The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. <a href="https://hypothesis.2">hypothesis.2</a> . While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. <a href="https://hypothesis.3">hypothesis.3</a> . Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON	Access of the Dalling Bridge	L C 1 L - P 26 - 1			
14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND	Approved for Public Release; Distri	Approved for Public Release; Distribution Unlimited			
14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND					
14. ABSTRACT This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND					
This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND	13. SUPPLEMENTARY NOTES				
This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND					
This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND					
This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. The study population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND	14 ADSTDACT				
population is male OEF/OIF/OND veterans who will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON		, of dolivering prolong	ad avacques for the	traatmant of F	OTCD in a group format. The study
Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON					
processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. <a href="https://www.hypothesis.1">hypothesis.1</a> . The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. <a href="https://www.hypothesis.2">hypothesis.2</a> . While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. <a href="https://www.hypothesis.3">hypothesis.3</a> . Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS  PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON					
longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. <a href="https://www.hypothesis1">https://www.hypothesis1</a> . The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. <a href="https://www.hypothesis2">https://www.hypothesis2</a> . While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. <a href="https://www.hypothesis3">https://www.hypothesis3</a> . Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON					
for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. <a href="Hypothesis1">Hypothesis 1</a> . The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. <a href="Hypothesis 2">Hypothesis 2</a> . While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. <a href="Hypothesis 3">Hypothesis 3</a> . Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS  PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON					
psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. <a href="Hypothesis1">Hypothesis1</a> . The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. <a href="Hypothesis2">Hypothesis 2</a> . While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. <a href="Hypothesis3">Hypothesis 3</a> . Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS  PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON	longer result in anxious arousal to t	rauma followed by esc	cape or avoidance b	ehaviors. PC1	is a non-trauma focused treatment
psycho-education regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. <a href="Hypothesis1">Hypothesis1</a> . The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. <a href="Hypothesis2">Hypothesis 2</a> . While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. <a href="Hypothesis3">Hypothesis 3</a> . Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS  PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON	for PTSD, where the mechanisms of	of change include alter	ing current maladap	tive relational	patterns and behaviors, providing
use of problem solving strategies to address present life problems. Hypothesis 1. The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2. While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON					
significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. <a href="Hypothesis 2">Hypothesis 2</a> . While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. <a href="Hypothesis 3">Hypothesis 3</a> . Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON					
both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON					
compared to the PCT group. Hypothesis 3. Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON					
will significantly decrease for subjects in the PEG compared with subjects in the PCT group.  15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON					
15. SUBJECT TERMS PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF: 17. LIMITATION 18. NUMBER 19a. NAME OF RESPONSIBLE PERSON					
PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON	will significantly decrease for subjects in the PEG compared with subjects in the PCT group.			).	
PTSD, Group Treatment, Male Veterans, OEF/OIF/OND  16. SECURITY CLASSIFICATION OF:  17. LIMITATION  18. NUMBER  19a. NAME OF RESPONSIBLE PERSON	15. SUBJECT TERMS				
16. SECURITY CLASSIFICATION OF: 17. LIMITATION 18. NUMBER 19a. NAME OF RESPONSIBLE PERSON		erans OFF/OIF/OND			
	40.050110177.01.4001710.1710.15		4= 1 1541= 4=	40 1111111	10 NAME OF BEOCH 10101 - 5-5-5-11
	To. SECURITY CLASSIFICATION OF:				

Unclassified

19b. TELEPHONE NUMBER (include area

code)

# **Table of Contents**

# **Page**

1. Introduction	2
2. Keywords	2
3. Accomplishments	2
4. Impact	5
5. Changes/Problems	5
5. Products	6
7. Participants & Other Collaborating Organizations	6
3. Special Reporting Requirements	10
O. Appendices	10

- 1. INTRODUCTION: The efficacy of exposure therapy has been established when delivered in an individual format. This study will establish the efficacy of delivering prolonged exposure for the treatment of PTSD in a group format. We will recruit and assess 376 male OEF/OIF/OND veterans, and with an estimated 20% dropout rate, 300 subjects will be block randomized in groups of 3 into a 10-week, 90-minute, Prolonged Exposure Group (PEG) or Present-Centered Therapy (PCT) group. The goal of the PEG therapy is to promote emotional processing of the trauma memory and desensitize anxiety reactions to trauma memories, so that memories or situations no longer result in anxious arousal to trauma followed by escape or avoidance behaviors. PCT is a non-trauma focused treatment for PTSD, where the mechanisms of change include altering current maladaptive relational patterns and behaviors, providing psychoeducation regarding the relationship between trauma and current relational patterns and behaviors, and teaching the use of problem solving strategies to address present life problems. Hypothesis 1: The Prolonged Exposure Group treatment will significantly lower PTSD symptoms compared to the Present-Centered Therapy control group. Hypothesis 2: While subjects in both groups will improve in self-reported overall PTSD symptomatology, the PEG will show greater within group improvement compared to the PCT group. Hypothesis 3: Perceived stigma associated with having a PTSD diagnosis and seeking treatment will significantly decrease for subjects in the PEG compared with subjects in the PCT group.
- 2. **KEYWORDS:** PTSD, Group Treatment, Male Veterans, OEF/OIF/OND.

#### 3. ACCOMPLISHMENTS:

• What were the major goals of the project?

Major Task 1: Adapt the Prolonged Exposure (PE) protocol to be delivered in a group format (months 1 through 6).

Major Task 2: Recruiting, Hiring, and Training Study Staff, and Securing Space Allocation for New Staff (months 1 through 6).

Major Task 3: Subject Recruitment, Baseline Assessment, and Randomization (months 7 through 33).

Major Task 4: Post-Treatment, 3– and 6-Month Follow-up Assessments (months 6 through 39)

Major Task 5: Data Entry (weekly), Reliability (quarterly) and Fidelity Checks (yearly) Major Task 6: Disseminating Findings at Professional Conference Presentations, and through Abstracts, Publications, and DoD (months 33 through 48)

a. What was accomplished under these goals?

## Major Task 1

- 1. Adapted individual Prolonged Exposure protocol for group delivery
- 2. Prepared Regulatory Documents and Research Protocol
- 3. Refined eligibility criteria, exclusion criteria, screening protocol
- 4. Finalized consent form and human subjects protocol

- 5. Completed and submitted IRB paperwork
- 6. Obtained IRB approval on 01/13/2015
- 7. Obtained R&D approval on 03/31/2015
- 8. Finalized assessment measurements and obtained instruments
- 9. Submitted amendments as needed, approval on 09/09/2015
- 10. Obtained DoD approval on 09/28/2015

### Major Task 2

- 1. Developed job descriptions
- 2. Advertised and interviewed potential staff
- 3. Selected staff, offered positions
- 4. Secured offices for new staff
- 5. Hired Psych Tech, RA, Therapists by 08/2015
- 6. Standardized training of staff on study protocol:
  - Therapists: Prolonged Exposure and Present-Centered Therapy protocols
  - Psych Tech/RA: Assessment Instruments, Motivational techniques, database development/management
- 7. Developed database and data entry screens

## Major Task 3

- 1. Present study to NMVAHCS Men's PTSD clinic staff, provide brochures/flyers
- 2. Present study to NMVAHCS OEF/OIF/OND Coordinator, provide brochures/flyers
- 3. Contact CBOCs and Vet Center to present study to clinic staff, provide brochures/flyers

#### **Goals not met:**

Several potential Psychologists/Study Coordinators applied, and some were qualified and interviewed; offers were made to four applicants over the past year, however, the position remained open. The most frequently cited reason was the desire to obtain a permanent position within the VA system. The position for the Psychologist/Study Coordinator has recently been accepted by a well-qualified applicant, and this person is going through the hiring process now.

Recruitment could not begin until DoD and IRB approvals were obtained. The DoD approval was on 09/28/2015, and we began recruitment efforts there afterward.

# b. What opportunities for training and professional development has the project provided?

- 1. All study personnel have been trained in-house on the study protocol.
- 2. The therapist who will be conducting the group Prolonged Exposure was trained by Dr Diane Castillo, and is working toward certification.
- 3. The group Present-Centered Therapy protocol was modified for this study in consultation with Dr Tracie Shea, and included the study therapist.
- 4. The Psych Tech and Research Assistant and other study staff attended CAPS-5 training conducted by Dr Frank Weathers.

## c. How were the results disseminated to communities of interest?

Nothing to Report

## d. What do you plan to do during the next reporting period to accomplish the goals?

The next reporting period will focus on recruitment:

Subject Recruitment, Baseline Assessments, and Randomization

Post-Treatment, 3– and 6-Month Follow-up Assessments

Data Entry, Reliability and Fidelity Checks

## 4. IMPACT:

a. What was the impact on the development of the principal discipline(s) of the project?

Nothing to Report

b. What was the impact on other disciplines?

Nothing to Report

c. What was the impact on technology transfer?

Nothing to Report

d. What was the impact on society beyond science and technology?

Nothing to Report

## 5. CHANGES/PROBLEMS:

a. Changes in approach and reasons for change

Nothing to Report

b. Actual or anticipated problems or delays and actions or plans to resolve them

The problem of hiring a Study Coordinator has been resolved, and Dr. Tim Ozechowski has been hired.

The delay in recruitment is resolved, and we are currently actively recruiting subjects.

c. Changes that had a significant impact on expenditures

Nothing to Report

d. Significant changes in use or care of human subjects, vertebrate animals, biohazards, and/or select agents

Nothing to Report

e. Significant changes in use or care of human subjects

Nothing to Report

- f. Significant changes in use or care of vertebrate animals. N/A
- g. Significant changes in use of biohazards and/or select agents. N/A
- 6. PRODUCTS: N/A

# 7. PARTICIPANTS & OTHER COLLABORATING ORGANIZATIONS

a. What individuals have worked on the project?

Name:	K. Janet C' de Baca
Project Role:	PI
Researcher Identifier (e.g. ORCID ID):	
Nearest person month worked:	4.8
Contribution to Project:	The PI is responsible for the administration and coordination of the grant and supervision of all study personnel; supervising the recruitment, testing, and treatment of all study participants, and ensuring the test and treatment data/results are reasonable, entered correctly, and managed appropriately to ensure accuracy and to protect patient confidentiality; and assuring the analysis of the data is performed with integrity, and preparing results for presentation and publication in peer-reviewed journals.
Funding Support:	

Name:	Diane Castillo
Project Role:	Co-PI
Researcher Identifier (e.g.	

ORCID ID):	
Nearest person month worked:	0.60
Contribution to Project:	The Co-PI will be able to step-in as needed for the PI in administering the grant and supervision of study personnel, and will assist in the preparation of results for presentation and publication.
Funding Support:	

Name:	Catherine Hearne
Project Role:	Co-Investigator
Researcher Identifier (e.g. ORCID ID):	
Nearest person month worked:	0.60
Contribution to Project:	The Co-Investigator, who is the OEF/OIF/OND point-of-contact and works primarily with recently returned Veterans, facilitating transition with the OEF/OIF/OND team will work with the research team to facilitate integration of the proposed project into her clinic, develop effective referral procedures, participate in resolving obstacles to recruitment, and participate in data analysis and manuscript preparation.
Funding Support:	

Name:	Christine Chee
Project Role:	Research Monitor
Researcher Identifier (e.g. ORCID ID):	
Nearest person month worked:	0.60
Contribution to Project:	The Research Monitor is responsible to oversee the safety of the research and report observations/findings to the NMVAHCS IRB or a designated official, and will review all unanticipated problems involving risks to subjects or others associated with the protocol and provide an independent report of the event to the NMVAHCS IRB.

	The Research Monitor may discuss the research protocol with the investigators, shall have authority to stop the research protocol in progress, remove individual human subjects from the study, and take whatever steps are necessary to protect the safety and well-being of human subjects until the IRB can assess the monitor's report; and shall have the responsibility to promptly report their observations and findings to the NMVAHCS IRB or other designated official and the HRPO.
Funding Support:	

Name:	Jenna Keller
Project Role:	Psychology Technician
Researcher Identifier (e.g. ORCID ID):	
Nearest person month worked:	12.0
Contribution to Project:	The Psychology Technician is responsible for performing assessments on study subjects (baseline, post-treatment, 3- and 6-month follow-up assessments); administering self-report instruments; and assisting with recruitment, scheduling, and data entry as needed.
Funding Support:	

Name:	Nichole Mays
Project Role:	Research Assistant
Researcher Identifier (e.g. ORCID ID):	
Nearest person month worked:	12.00
Contribution to Project:	The Research Assistant is responsible for assisting with assessments, scheduling and tracking post-treatment, 3- and 6-month follow-up assessments in a timely manner to assure study guidelines; and entering data accurately; and assisting with the recruitment of subjects.

Funding Support:	
------------------	--

Name:	Elizabeth McCallion
Project Role:	Therapist
Researcher Identifier (e.g. ORCID ID):	
Nearest person month worked:	6.00
Contribution to Project:	This Master's level clinician has been trained in the group Prolonged Exposure therapy and will be responsible for conducting 50, 3-person, 90-minute, 10-session group Prolonged Exposure therapy sessions over a 33 month period; and for maintaining subject treatment files, administering/scoring PCL-5's, monitoring session audiorecordings by subjects, providing handouts, and videotaping sessions for fidelity checks.
Funding Support:	

Name:	Elizabeth McLaughlin
Project Role:	Therapist
Researcher Identifier (e.g. ORCID ID):	
Nearest person month worked:	6.00
Contribution to Project:	This Master's level clinician has been trained in the group Present-Centered Therapy protocol and will be responsible for conducting 50, 3-person, 90-minute, 10-session Present-Centered Therapy groups over a 33 month period; and for maintaining subject treatment files, administering/scoring PCL-5's, providing handouts, and videotaping sessions for fidelity checks.
Funding Support:	

b. Has there been a change in the active other support of the PD/PI(s) or senior/key personnel since the last reporting period?

Nothing to Report

- c. What other organizations were involved as partners? N/A
- 8. SPECIAL REPORTING REQUIREMENTS N/A
- 9. APPENDICES: N/A

DO NOT RENUMBER PAGES IN THE APPENDICES.